

On The Arnold Conjecture And The Atiyah-Patodi-Singer Index Theorem

Mauri Miettinen^{*} and Antti J. Niemi[#]

^{}# Department of Theoretical Physics, Uppsala University, P.O. Box 803,
S-75108, Uppsala, Sweden*

^{}# The Mittag-Leffler Institute, Auravägen 17, S-182 62 Djursholm, Sweden*

*[#] Helsinki Institute of Physics, P.O. Box 9, FIN-00014 Helsinki University,
Finland*

The Arnold conjecture yields a lower bound to the number of periodic classical trajectories in a Hamiltonian system. Here we count these trajectories with the help of a path integral, which we inspect using properties of the spectral flow of a Dirac operator in the background of a $\mathrm{Sp}(2N)$ valued gauge field. We compute the spectral flow from the Atiyah-Patodi-Singer index theorem, and apply the results to evaluate the path integral using localization methods. In this manner we find a lower bound to the number of periodic classical trajectories which is consistent with the Arnold conjecture.

[#] niemi@teorfys.uu.se

Periodic classical trajectories are important in a large number of problems. Examples range from classical integrable models where the motion is a periodic flow on invariant torii, to quantum chaos where the energy spectrum is computed by the Gutzwiller (Selberg) trace formula. For non-integrable models exact analytical results are sparse, and in general even a classification of trajectories becomes arduous. Some 35 years ago Arnold conjectured [1] that in an arbitrary Hamiltonian system the number of periodic classical solutions is bounded from below by the number of critical points of a function. If the Hamiltonian is energy conserving *i.e.* has no explicit time dependence, this follows immediately from conventional Morse inequalities since every critical point of the Hamiltonian is trivially a periodic classical solution. But in the general case when the Hamiltonian is explicitly time dependent so that energy is not conserved, the conjecture has only been verified in special cases. The modern approach to the Arnold conjecture is due to Floer [2] who formulated it using infinite dimensional Morse theory. His approach has led to extensive investigations [3] and applications in seemingly disparate subjects, most notably topological quantum field theories and strings [4].

Here we shall study relations between the Arnold conjecture and canonical path integrals. We suggest an intuitive approach for proving the conjecture using path integral localization techniques. Our arguments are based on methods of topological quantum field theories and properties of spectral flow in a 1+1 dimensional Dirac operator in the background of an adiabatically varying $\text{Sp}(2N)$ valued classical gauge field. For this we consider Hamilton's equation of motion on a $2N$ -dimensional phase space *i.e.* a symplectic manifold Γ with local coordinates x^i ,

$$\dot{x}^i = \omega^{ij} \partial_j H(x, t) \equiv \mathcal{X}_H^i(x, t) \quad (1)$$

Here $H(x, t)$ is in general an explicitly time dependent Hamiltonian and $\mathcal{X}_H(x, t)$ is the corresponding Hamiltonian vector field. The antisymmetric $\omega^{ij} = \{x^i, x^j\}$ is the matrix inverse of the symplectic two-form $\omega = \frac{1}{2} \omega_{ij} dx^i \wedge dx^j$ on Γ . The solutions of (1) are critical points of the classical action

$$S = \int dt \left[\vartheta_i \dot{x}^i - H(x, t) \right] \quad (2)$$

where ϑ is the symplectic potential, $\omega_{ij} = \partial_i \vartheta_j - \partial_j \vartheta_i$. For a T -periodic Hamiltonian $H(x, t) = H(x, t + T)$ defined on a compact Γ the Arnold conjecture states that the number of non-degenerate and contractible T -periodic solutions is bounded from below by the sum of the Betti numbers B_p on Γ ,

$$\#\{\text{T-periodic solutions}\} \geq \sum B_p \quad (3)$$

When the t -dependence of H is trivial (3) reduces to the conventional finite dimensional Morse inequality [5] since every critical point of H is trivially T -periodic. But for a generic t -dependent Hamiltonian (3) remains unproven.

The number of T -periodic solutions is counted by the path integral

$$Z = \sum_{\delta S=0} 1 = \int \mathcal{D}x \, \delta \left[\frac{\delta S}{\delta x} \right] \cdot \left| \text{Det} \left[\frac{\delta^2 S}{\delta x^2} \right] \right| \quad (4)$$

where we have the determinant of the Hessian

$$\frac{\delta^2 S}{\delta x^i(t_1) \delta x^j(t_2)} = \left\{ \omega_{ij} \partial_{t_1} + \frac{1}{2} (\partial_j \omega_{ik} + \partial_i \omega_{jk}) \dot{x}^k - \partial_i \partial_j H \right\} \delta(t_1 - t_2) \quad (5)$$

which is a symmetric first-order differential operator. The zero-modes of (5) are Jacobi fields, but we assume that the T -periodic solutions of (1) are isolated and non-degenerate so that the Jacobi fields are absent. The difficulty in establishing (3) is then due to the absolute signs in (4), alternatively due to the first-order nature of the Hessian: The Morse index of (2) coincides with the number of negative eigenvalues of (5). But since the spectrum of (5) is unbounded from below this number is infinite and needs to be regulated appropriately.

We first consider (4) without the absolute value. Formally, the path integral then evaluates to the Euler characteristic on the loop space $\text{L}\Gamma$,

$$Z = \sum_{\delta S=0} (-)^k = \int \mathcal{D}x \, \delta \left[\frac{\delta S}{\delta x} \right] \text{Det} \left[\frac{\delta^2 S}{\delta x^2} \right] \quad (6)$$

where $k = 0$ or 1 depending on whether the Morse index of the pertinent critical point of S is even or odd, *i.e.* whether the Hessian has an even or odd number of negative eigenvalues in some proper regularization scheme.

We introduce a commuting Lagrange multiplier λ and anticommuting ghosts $\psi, \bar{\psi}$, and re-write (6) as

$$\begin{aligned} Z &= \int \mathcal{D}\Phi \exp \left\{ i \int_0^T dt \left[\lambda_i (\dot{x}^i - \mathcal{X}_H^i) + \bar{\psi}_i (\delta_j^i \partial_t + \partial_j \mathcal{X}_H^i) \psi^j \right] \right\} \\ &= \int \mathcal{D}\Phi \exp \left\{ i \int_0^T dt \left[\lambda_i \dot{x}^i + \bar{\psi}_i \dot{\psi}^i - \lambda_i \mathcal{X}_H^i - \psi^i \partial_i \mathcal{X}_H^j \bar{\psi}_j \right] \right\} \end{aligned} \quad (7)$$

where we denote by Φ collectively the T -periodic variables $(x, \lambda, \psi, \bar{\psi})$. This has the structure of a canonical path integral in a graded phase space in Darboux coordinates with Poisson brackets

$$\{\lambda_i, x^j\} = \{\bar{\psi}_i, \psi^j\} = \delta_i^j \quad (8)$$

We note that the Lagrange multiplier λ provides a realization of derivation *w.r.t.* the variable x . Similarly we identify $\psi \sim dx$, the basis one-forms, and $\bar{\psi}$ is the corresponding basis of contractions. We define the exterior derivative in the loop space $L\Gamma$

$$d = \int_0^T dt \psi^i \lambda_i \quad (9)$$

We may view d as a nilpotent BRST operator and the action S_{ext} in (7) is a BRST commutator,

$$S_{\text{ext}} = \lambda_i \dot{x}^i + \bar{\psi}_i \dot{\psi}^i - \lambda_i \mathcal{X}_H^i - \psi^i \partial_i \mathcal{X}_H^j \bar{\psi}_j = [d, \Psi] = \left[d, (\dot{x}^i - \mathcal{X}_H^i) \bar{\psi}_i \right] = \mathcal{L}_S \quad (10)$$

with $\mathcal{L}_S$ the Lie derivative along the loop space vector field $\mathcal{X}_S = \dot{x} - \mathcal{X}_H$. These relations mean that the path integral (7) determines a topological quantum theory [6].

The Fradkin-Vilkovisky theorem [7] states that (7) is invariant under an infinitesimal variation $\Psi \rightarrow \Psi + \delta\Psi$ of the gauge fermion in (10). An arbitrary infinitesimal coordinate transformation $x \rightarrow \tilde{x} = x + \epsilon(x)$ in the original phase space Γ is an example of such a variation, with

$$\Psi \rightarrow \Psi + \frac{\partial \epsilon^i}{\partial x^j} (\dot{x}^j - \mathcal{X}_H^j) \bar{\psi}_i$$

provided we also conjugate the BRST operator according to

$$d \rightarrow e^{-\lambda_i \epsilon^i} d e^{\lambda_i \epsilon^i}$$

Since this conjugation does not alter the cohomology of d it has no effect on the path integral, which is left intact. As a consequence we can inspect the local structure of the path integral by selecting (local) Darboux coordinates on Γ ,

$$\omega \rightarrow \omega_0 = \sum_{k=1}^N dp_k \wedge dq^k \quad (11)$$

In these coordinates

$$A^j_i = \partial_i \mathcal{X}^j_H = \omega_0^{jk} \partial_i \partial_k H \quad (12)$$

is a $\text{Sp}(2N)$ -valued traceless product of the antisymmetric ω_0^{-1} and the symmetric Hessian $\partial^2 H$, and the ghost part in (7) simplifies into

$$S_F = \int_0^T dt \bar{\psi}(\partial_t + A)\psi \quad (13)$$

This is reminiscent of a (0+1) dimensional Dirac Lagrangian in the background of a gauge field. Indeed, the Darboux coordinates are defined only modulo canonical transformations corresponding to arbitrary t -dependent conjugations of the symplectic two-form (11) by elements of $\text{Sp}(2N)$,

$$\omega_0 = U^{-1}(t)\omega_0 U(t)$$

Under these $\text{Sp}(2N)$ conjugations the anticommuting ghosts transform according to

$$\begin{aligned} \psi(t) &\rightarrow U(t)\psi(t) \\ \bar{\psi}(t) &\rightarrow \bar{\psi}(t)U^{-1}(t) \end{aligned} \quad (14)$$

and (13) remains invariant if $A^j_i(t)$ transforms like a $\text{Sp}(2N)$ -valued gauge field

$$A(t) \rightarrow U(t)A(t)U^{-1}(t) + U(t)\dot{U}^{-1}(t) \quad (15)$$

Following [8] we introduce

$$\Lambda(t) = \text{T exp} \left(\int_0^t A(t') dt' \right)$$

and define a $\text{Sp}(2N)$ gauge transformation with

$$U(t) = e^{-A_0 t} \cdot \Lambda^\dagger(t) \quad (16)$$

where A_0 is defined by

$$\Lambda(T) = \exp(-A_0 \cdot T) \quad (17)$$

which is essentially the average of $A(t)$ over the period T . This gauge transformation maps our $\text{Sp}(2N)$ -valued gauge field $A(t)$ into the t -independent average A_0 ,

$$U(t)A(t)U^{-1}(t) + U(t)\dot{U}^{-1}(t) = A_0 \quad (18)$$

As a consequence the ghost Lagrangian (13) depends only on the t -independent A_0 . Moreover, if we decompose the T -periodic Hamiltonian $H(x, t)$ into its Fourier modes

$$H(x, t) = H_0(x) + \sum_{n \neq 0} H_n(x) \cdot \exp\left(\frac{2\pi i n t}{T}\right) \quad (19)$$

where the $H_n(x)$ have no explicit t -dependence and $H_0(x)$ is the average of $H(x, t)$ over the period T , we conclude that the gauge transformation (18) replaces the original Hamiltonian $H(x, t)$ by its t -independent average $H_0(x)$ in the ghost Lagrangian (13).

In the following we shall assume that the average $H_0(x)$ is a Morse function with only isolated and non-degenerate critical points.

Let Ψ denote the original gauge fermion in (10) and let Ψ_0 denote the corresponding gauge fermion with $H(x, t)$ replaced by the average $H_0(x)$. We define the infinitesimal change of variables

$$\Phi \rightarrow \Phi + \epsilon \cdot \{d, \Phi\} \cdot \int_0^T (\Psi - \Psi_0) \quad (20)$$

with ϵ an infinitesimal parameter. We find that in (7), (10) the *only* effect of (20) is the shift

$$\Psi \rightarrow (1 - \epsilon)\Psi + \epsilon\Psi_0$$

and by repeating this change of variables we can replace $H(x, t)$ by $H_0(x)$ in (7). Hence (7) depends only on the t -independent average $H_0(x)$ of the Hamiltonian.

The Hessian (5) has an eigenvalue problem

$$Df_n(t) = \lambda_n f_n(t) \quad , \quad f_n(0) = f_n(T) \quad (21)$$

where the eigenvalues λ_n are invariant under $\text{Sp}(2N)$ gauge transformations that are continuously connected to the identity. Since we assume that there are no Jacobi fields, in the background of a classical solution all $\lambda_n \neq 0$. If we introduce the η -invariant of the Hessian

$$\eta_D = \lim_{s \rightarrow 0} \sum_{\lambda_n \neq 0} \text{sign}(\lambda_n) |\lambda_n|^{-s} \quad (22)$$

we can remove the absolute value of the determinant in (4) by

$$\left| \text{Det}[D] \right| \rightarrow \text{Det}[D] \cdot \exp\left(\frac{i\pi}{2}\eta_D\right) \quad (23)$$

This implies that the number of T -periodic classical trajectories can be counted by the following generalization of (7),

$$\mathcal{Z} = \int \mathcal{D}\Phi \exp \left\{ i \int_0^T dt \left[\lambda_k \dot{x}^k + \bar{\psi}_k \dot{\psi} - \lambda_k \mathcal{X}_H^k - \psi^k \partial_k \mathcal{X}_H^l \bar{\psi}_l \right] + \frac{i\pi}{2}\eta_D \right\} \quad (24)$$

The path integral (7) depends only on the t -independent average $H_0(x)$. In particular, in the ghost action (13) the Hamiltonian $H(x, t)$ can be converted into $H_0(x)$ by a $\text{Sp}(2N)$ gauge transformation which leaves the η -invariant of the Hessian intact. As a consequence we conclude that the path integral (24) should not change if we replace $H(x, t) \rightarrow H_0(x)$ in it.

The Atiyah-Patodi-Singer index theorem [9] relates the η -invariant of a $(0+1)$ dimensional Dirac operator to the index of a $(1+1)$ dimensional

Dirac operator. Accordingly we extend our Hessian (5) to the following two-dimensional Dirac operator on a cylinder $S^1 \times \mathbf{R}$

$$\mathcal{Q} = \begin{pmatrix} 0 & \partial_s + D_s \\ -\partial_s + D_s & 0 \end{pmatrix} = i\sigma_2 \otimes \partial_s + \sigma_1 \otimes D_s = \begin{pmatrix} 0 & Q^\dagger \\ Q & 0 \end{pmatrix} \quad (25)$$

Here D_s with $s \in \mathbf{R}$ is a one-parameter family of symmetric Hessians on the cylinder. We define them by promoting our T -periodic solution $x(t)$ into a one-parameter family of T -periodic trajectories $u(s, t)$, an instanton that adiabatically interpolates between $x(t)$ at $s \rightarrow +\infty$ and another T -periodic $y(t)$ at $s \rightarrow -\infty$,

$$\begin{aligned} \lim_{s \rightarrow +\infty} u(s, t) &= x(t) \\ \lim_{s \rightarrow -\infty} u(s, t) &= y(t) \end{aligned} \quad (26)$$

The operators D_s have an s -dependent eigenvalue problem

$$D_s f_n(t, s) = \lambda_n(s) f_n(t, s) \quad , \quad f_n(0, s) = f_n(T, s) \quad (27)$$

and in the absence of Jacobi fields at $s \rightarrow \pm\infty$ all $\lambda_n(\pm\infty) \neq 0$. But as a function of the adiabatic parameter s , the operators D_s in general exhibit a non-trivial spectral flow with some of the eigenvalues $\lambda_n(s)$ crossing zero when the parameter s evolves. This spectral flow can be computed by applying the Atiyah-Patodi-Singer index theorem to the Dirac operator $\mathcal{Q}$. It relates the index of $\mathcal{Q}$ on the cylinder to the η -invariants of the operator D_s at the $s \rightarrow \pm\infty$ ends of the cylinder,

$$\begin{aligned} \text{ind } \mathcal{Q} &= \dim \text{Ker } Q - \dim \text{Ker } Q^\dagger \\ &= \frac{1}{2} (\eta_D[y(t)] - \eta_D[x(t)]) + \frac{1}{2\pi} \int_{S^1 \times \mathbf{R}} \text{tr } F \end{aligned} \quad (28)$$

Here $F = \partial_s(\omega_0 A)$ is a $\text{Sp}(2N)$ -valued field strength: We have a two dimensional $\text{Sp}(2N)$ gauge field A with components $A_t = \omega_0 A$ and $A_s = 0$. In general the contribution from F cancels the continuous part in the difference of the η -invariants in (28). But since we have verified that in the Darboux variables (11) the $\text{Sp}(2N)$ valued gauge field $\omega_0 A$ is explicitly traceless, the relative η -invariant in (28)

$$\Delta\eta_D = \eta_D[y(t)] - \eta_D[x(t)] \quad (29)$$

should be an integer, it coincides with the net spectral flow between the two configurations $x(t)$ and $y(t)$,

$$\Delta\eta_D \sim \text{spectral flow of } \mathcal{Q} \quad (30)$$

In particular (29) determines the relative sign between the determinants of the Hessians for the two trajectories $y(t)$ and $x(t)$. Hence $\Delta\eta_D$ coincides with the relative Morse index between the two trajectories, which is also the difference between their Maslov indices [3].

Instead of (24) we proceed with the more general

$$\mathcal{Z} = \int \mathcal{D}\Phi \exp \left[i \int_0^T dt \left(\lambda_k \dot{x}^k + \bar{\psi}_k \dot{\psi} - \lambda_k \mathcal{X}_H^k - \psi^k \partial_k \mathcal{X}_H^l \bar{\psi}_l \right) + \frac{i\pi}{2} \Delta\eta_D \right] . \quad (31)$$

with $\Delta\eta_D$ the relative η -invariant *w.r.t.* a fixed T -periodic reference trajectory $y(t)$. It is integer valued, and twice the index of $\mathcal{Q}$ in the background of the pertinent instanton $u(s, t)$. Since (24) remains intact when we replace $H(x, t)$ by its average $H_0(x)$, we can expect that (31) also remains intact when we set $H(x, t) \rightarrow H_0(x)$.

Since the relative η -invariant in (31) coincides with the integer valued index of the Dirac operator $\mathcal{Q}$, it should not change under a continuous change of variables. This suggests that we can evaluate the integral (31) by localization. Accordingly we decompose the variables Φ in their t -independent and t -dependent components, $\Phi \rightarrow \Phi_0 + \Phi_t$, and set $x(t) \rightarrow x_0 + \alpha x_t$, $\psi(t) \rightarrow \psi_0 + \sqrt{\alpha} \psi_t$, and $\bar{\psi}(t) \rightarrow \bar{\psi}_0 + \sqrt{\alpha} \bar{\psi}_t$ with α a parameter. The corresponding super-Jacobian in (31) is trivial, and we localize by setting $\alpha \rightarrow 0$. In this limit the path integral (7) is known to localize to the Poincaré-Hopf representation of the Euler characteristic of the phase space Γ [10],

$$Z = \sum_{\mathcal{X}_{H_0}=0} \text{sign} \det(\partial \mathcal{X}_{H_0}) = \sum_{k=0}^{2N} (-1)^k m_k = \sum_{k=0}^{2N} (-1)^k B_k = \chi(\Gamma) \quad (32)$$

where m_k denotes the number of critical points with Morse index k . In a similar manner we expect the path integral (31) to localize into

$$\mathcal{Z} = \sum_{\mathcal{X}_{H_0}=0} \text{sign} \det(\partial \mathcal{X}_{H_0}) \exp \left(+ \frac{i\pi}{2} \Delta\eta_D \right) \quad (33)$$

where $\Delta\eta_D$ is now the relative η -invariant for the critical points of $H_0(x)$ *w.r.t.* the fixed periodic reference trajectory $y(t)$. We select this reference trajectory so that it coincides with a critical point of $H_0(x)$ with Morse index 0. According to the standard finite dimensional Morse inequality $m_p \geq B_p$ and $B_0 \geq 1$ which ensures that such a critical point always exists. With this, the pertinent Dirac index (28) simplifies into

$$\text{ind } Q = \dim \text{Ker } Q - \dim \text{Ker } Q^\dagger = \dim \text{Ker } Q \quad (34)$$

This follows directly from the zero mode equation in the adiabatic limit, where we can use the Ansatz

$$g_n(s, t) = \phi_n(s) \rho_n(t)$$

with

$$D_s \rho_n(t) = \lambda_n(s) \rho_n(t)$$

This reduces the zero mode equations for Q and $Q^\dagger$ to

$$(\partial_s \pm \lambda_n(s)) \phi_n(s) = 0$$

with solutions

$$\phi_n(s) = \exp \left[\mp \int_0^s ds' \lambda_n(s') \right] \phi_n(0)$$

respectively. Hence $Q^\dagger = \partial_s + D_s$ does not have a normalizable zero mode and (34) follows. In particular, the spectral flow is only from positive eigenvalues of the Hessian to its negative eigenvalues, which implies that the contribution from the relative η -invariant $\Delta\eta_D$ in (33) coincides with the Morse index $m_{H_0}(x) = k$ of the critical point. Since $\text{sign } \det(\partial\mathcal{X}_{H_0}) = (-1)^k$ and Morse inequalities yield $m_k \geq B_k$, we then conclude that modulo an overall, irrelevant sign the path integral (31) localizes into

$$Z = \sum_{\mathcal{X}_{H_0}=0} \text{sign } \det(\partial\mathcal{X}_{H_0}) \exp(-i\pi k) = \sum_{\mathcal{X}_{H_0}=0} 1 = \sum_{k=0}^{2N} m_k \geq \sum_{k=0}^{2N} B_k$$

As a consequence we have a lower bound estimate for the number of periodic classical trajectories which is consistent with the Arnold conjecture.

In conclusion, we have studied relations between a path integral and the Arnold conjecture. We have related the path integral to the properties of a two dimensional Dirac operator and applied the Atiyah-Patodi-Singer index theorem to evaluate it using localization methods. In this manner we have found a lower bound estimate for the number of periodic classical trajectories which is consistent with the Arnold conjecture. Our approach is based on intuitive path integral arguments, and we hope that a rigorous derivation of the conjecture could be developed along these lines.

We thank Topi Kärki for discussions. Our research has been partially supported by NFR Grant F-AA/FU 06821-308. This article reports collaboration that was largely completed by March 2nd 1999, when Mauri Miettinen unexpectedly passed away just one month before his scheduled Thesis defence.

References

- [1] V.I. Arnold, C.R. Acad. Paris **261** (1965) 3719; and Usp. Math. Nauk. **18** (1963) 91
- [2] A. Floer, Comm. Pure Appl. Math. **41** (1988) 775, and *ibid.* **42** (1989) 775
- [3] A. Hofer and E. Zehnder, *Symplectic Invariants and Hamiltonian Dynamics* (Birkhäuser Verlag, 1994)
- [4] E. Witten, Comm. Math. Phys. **118** (1988) 411
- [5] J. Milnor, *Morse Theory* (Princeton University Press, 1973)
- [6] D. Birmingham, M. Blau, M. Rakowski and G. Thompson, Phys. Repts. **209** (1991) 129

- [7] E.S. Fradkin and G.A. Vilkovisky, Phys. Lett. **55** (1975) 224; for a review, see A.J. Niemi, Phys. Repts. **184** (1989) 147
- [8] Contribution by R. Jackiw in S.B. Treiman, E. Witten, R. Jackiw and B. Zumino, *Current Algebra And Anomalies* (World Scientific, Singapore 1985); S. Elitzur, E. Rabinovici, Y. Frishman and A. Schwimmer, Nucl. Phys. **B273** (1986) 93
- [9] M.F. Atiyah, V.K. Patodi and I.M. Singer, Math. Proc. Camb. Phil. Soc. **77** (1975) 43
- [10] A.J. Niemi, Phys. Lett. **B355** (1995) 501